

*Citation for published version:*

Vincze, O, Kosztolányi, A, Barta, Z, Küpper, C, Alrashidi, M, Amat, JA, Argüelles Ticó, A, Burns, F, Cavitt, J, Conway, WC, Cruz-López, M, Desucre-Medrano, AE, dos Remedios, N, Figuerola, J, Galindo-Espinosa, D, García-Peña, GE, Gómez Del Angel, S, Gratto-Trevor, C, Jönsson, P, Lloyd, P, Montalvo, T, Parra, JE, Pruner, R, Que, P, Liu, Y, Saalfeld, ST, Schulz, R, Serra, L, St Clair, JJH, Stenzel, LE, Weston, MA, Yasué, M, Zefania, S & Székely, T 2017, 'Parental cooperation in a changing climate: fluctuating environments predict shifts in care division', *Global Ecology and Biogeography*, vol. 26, no. 3, pp. 347-358. <https://doi.org/10.1111/geb.12540>

*DOI:*

[10.1111/geb.12540](https://doi.org/10.1111/geb.12540)

*Publication date:*

2017

*Document Version*

Peer reviewed version

[Link to publication](#)

This is the peer reviewed version of the following article: Orsolya Vincze András Kosztolányi Zoltán Barta Clemens Küpper Monif Alrashidi Juan A. Amat Araceli Argüelles Ticó Fiona Burns John Cavitt Warren C. Conway Medardo CruzLópez Atahualpa Eduardo DesucreMedrano Natalie dos Remedios Jordi Figuerola Daniel GalindoEspinosa Gabriel E. GarcíaPeña Salvador Gómez Del Angel Cheri GrattoTrevor Paul Jönsson Penn Lloyd Tomás Montalvo Jorge Enrique Parra Raya Pruner Pinjia Que Yang Liu Sarah T. Saalfeld Rainer Schulz Lorenzo Serra James J. H. St Clair Lynne E. Stenzel Michael A. Weston Maï Yasué Sama Zefania Tamás Székely (2016) Parental cooperation in a changing climate: fluctuating environments predict shifts in care division. *Global Ecology and Biogeography*, 26(3), which has been published in final form at [doi.org/10.1111/geb.12540](https://doi.org/10.1111/geb.12540). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



**Parental cooperation in a changing climate: fluctuating environments predict shifts in care division**

Journal:	<i>Global Ecology and Biogeography</i>
Manuscript ID	GEB-2016-0012
Manuscript Type:	Research Papers
Date Submitted by the Author:	10-Jan-2016
Complete List of Authors:	<p>Vincze, Orsolya; Debreceni Egyetem, Department of Evolutionary Zoology and Human Biology; Universitatea Babes-Bolyai, Hungarian Department of Biology and Ecology</p> <p>Kosztolányi, András; Debreceni Egyetem, Department of Evolutionary Zoology and Human Biology; Szent Istvan Egyetem, Department of Ecology</p> <p>Barta, Zoltán; Debreceni Egyetem, Department of Evolutionary Zoology and Human Biology</p> <p>Küpper, Clemens; University of Graz, Institute of Zoology</p> <p>AlRashidi, Monif; University of Ha'il, Department of Biology</p> <p>Amat, Juan; Estacion Biologica de Donana CSIC, Department of wetland Ecology</p> <p>Argüelles Ticó, Araceli; University of Bath, Department of Biology and Biochemistry</p> <p>Burns, Fiona; RSPB Centre for Conservation Science, RSPB Centre for Conservation Science</p> <p>Cavitt, John; Weber State University, Department of Zoology</p> <p>Conway, Warren; Texas Tech University, Department of Natural Resources Management</p> <p>Cruz-López, Medardo; Universidad Nacional Autónoma de México, Posgrado en Ciencias del Mar y Limnología</p> <p>Desucre-Medrano, Atahualpa Eduardo ; Universidad Nacional Autónoma de México, Posgrado en Ciencias del Mar y Limnología</p> <p>dos Remedios, Natalie; University of Sheffield, Department of Animal and Plant Sciences</p> <p>Figuerola, J; CSIC, Sevilla, Wetland Ecology</p> <p>Galindo-Espinosa, Daniel; Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas</p> <p>García-Peña, Gabriel E.; CESAB- Centre de Synthèse et d'Analyse sur la Biodiversité, CESAB- Centre de Synthèse et d'Analyse sur la Biodiversité; Centre de Recherche IRD -UMR MIVEGEC, Maladies Infectieuses et Vecteurs: Ecologie, Génétique, Evolution et Contrôle</p> <p>Del Angel, Salvador Gómez ; Universidad Nacional Autónoma de México, Posgrado en Ciencias del Mar y Limnología</p> <p>Gratto-Trevor, Cheri; Environment Canada, Prairie and Northern Wildlife Research Centre</p> <p>Jönsson, Paul; Lund University, Department of Biology</p> <p>Lloyd, Penn; University of Cape Town, Percy FitzPatrick Institute</p> <p>Montalvo, Tomás; Agència de Salut Pública de Barcelona, Servei de</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

	<p>Vigilància i Control de Plagues Urbanes Parra, Jorge Enrique; University of Bath, Department of Biology and Biochemistry Pruner, Raya; University of Florida, Department of Wildlife Ecology and Conservation Que, Pinjia; Beijing Normal University, Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences Liu, Yang; Sun Yat-sen University, State Key Laboratory of Biocontrol and College of Ecology and Evolution Saalfeld, Sarah; Manomet Center for Conservation Sciences, Manomet Center for Conservation Sciences Schulz, Rainer; Schutzstation Wattenmeer Nationalparkhaus, Schutzstation Wattenmeer Nationalparkhaus Serra, Lorenzo; Istituto Superiore per la Protezione e la Ricerca Ambientale, Istituto Superiore per la Protezione e la Ricerca Ambientale St Clair, James J.H.; University of St Andrews, School of Biology Stenzel, Lynne; Point Blue Conservation Science, Point Blue Conservation Science Weston, Michael; Deakin University, Centre for Integrative Ecology Yasué, Maï Yasué; Quest University Canada, Social Science Department Zefania, Sama; University of Toliara, Department of Biology Székely, Tamás ; University of Bath, Department of Biology and Biochemistry</p>
Keywords:	global change, climate, environmental stochasticity, seasonal environment, parental care, parental cooperation



# Parental cooperation in a changing climate: fluctuating environments predict shifts in care division

Orsolya Vincze<sup>1,2\*</sup>, András Kosztolányi<sup>1,3</sup>, Zoltán Barta<sup>1</sup>, Clemens Küpper<sup>4</sup>, Monif Alrashidi<sup>5</sup>, Juan A. Amat<sup>6</sup>, Araceli Argüelles Ticó<sup>7</sup>, Fiona Burns<sup>8</sup>, John Cavitt<sup>9</sup>, Warren C. Conway<sup>10</sup>, Medardo Cruz-López<sup>11</sup>, Atahualpa Eduardo Desucere-Medrano<sup>12</sup>, Natalie dos Remedios<sup>13</sup>, Jordi Figuerola<sup>6</sup>, Daniel Galindo-Espinosa<sup>14</sup>, Gabriel E. García-Peña<sup>15,16</sup>, Salvador Gómez Del Angel<sup>12</sup>, Cheri Gratto-Trevor<sup>17</sup>, Paul Jönsson<sup>18</sup>, Penn Lloyd<sup>19</sup>, Tomás Montalvo<sup>20</sup>, Jorge Enrique Parra<sup>7</sup>, Raya Pruner<sup>21</sup>, Pinjia Que<sup>22</sup>, Yang Liu<sup>23</sup>, Sarah T. Saalfeld<sup>24</sup>, Rainer Schulz<sup>25</sup>, Lorenzo Serra<sup>26</sup>, James J. H. St Clair<sup>27</sup>, Lynne E. Stenzel<sup>28</sup>, Michael A. Weston<sup>29</sup>, Mai Yasué<sup>30</sup>, Sama Zefania<sup>31</sup> and Tamás Székely<sup>7</sup>

<sup>1</sup>MTA-DE “Lendület” Behavioural Ecology Research Group, Department of Evolutionary Zoology and Human Biology, University of Debrecen, Debrecen, H-4032, Hungary

<sup>2</sup>Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj Napoca, RO-400006, Romania

<sup>3</sup>Department of Ecology, Faculty of Veterinary Science, Szent István University, Budapest, H-1077, Hungary

<sup>4</sup>Institute of Zoology, University of Graz, Universitätsplatz 2, 8010 Graz, Austria

<sup>5</sup>Department of Biology, Faculty of Science, University of Ha'il, Ha'il, PO Box 2440, Saudi Arabia

<sup>6</sup>Estación Biológica de Doñana (EBD-CSIC), calle Américo Vespucio s/n, Sevilla, 41092, Spain

<sup>7</sup>Biodiversity Lab, Department of Biology and Biochemistry, University of Bath, Bath, BA1 7AY, United Kingdom

<sup>8</sup>RSPB Centre for Conservation Science, RSPB, The Lodge, Sandy, Bedfordshire, SG192DL, United

1  
2 23 *Kingdom*  
3  
4 24 <sup>9</sup>*Avian Ecology Laboratory, Department of Zoology, Weber State University, Ogden, Utah, 84408, USA*  
5  
6 25 <sup>10</sup>*Department of Natural Resources Management, Texas Tech University, Lubbock, Texas, 79409, USA*  
7  
8 26 <sup>11</sup>*Posgrado en Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Ciudad*  
9  
10 *Universitaria, México D.F., 04510, México*  
11  
12 27  
13 28 <sup>12</sup>*Laboratorio de Zoología, Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma*  
14  
15 *de México, Tlalnepantla, 54000, México*  
16 29  
17 30 <sup>13</sup>*NERC Biomolecular Analysis Facility, Department of Animal and Plant Sciences, University of*  
18  
19 *Sheffield, Sheffield S10 2TN, UK*  
20  
21 31  
22 32 <sup>14</sup>*Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, Baja California Sur,*  
23  
24 *23096, México*  
25  
26 33  
27 34 <sup>15</sup>*CESAB- Centre de Synthèse et d'Analyse sur la Biodiversité, Aix-en-Provence cedex 3, 13857, France*  
28  
29 35 <sup>16</sup>*Centre de Recherche IRD -UMR MIVEGEC. Maladies Infectieuses et Vecteurs: Ecologie, Génétique,*  
30  
31 *Evolution et Contrôle, Montpellier cedex 5, 34394, France*  
32  
33 36  
34 37 <sup>17</sup>*Prairie and Northern Wildlife Research Centre, Science and Technology Branch, Environment*  
35  
36 *Canada, Saskatoon, SK S7N 0X4, Canada*  
37  
38 38  
39 39 <sup>18</sup>*Department of Biology, Lund University, Lund, S-22362, Sweden*  
40  
41 40 <sup>19</sup>*Percy FitzPatrick Institute, DST/NRF Centre of Excellence, University of Cape Town, Rondebosch,*  
42  
43 *7701, South Africa*  
44  
45 41  
46 42 <sup>20</sup>*Servei de Vigilància i Control de Plagues Urbanes, Agència de Salut Pública de Barcelona,*  
47  
48 *Barcelona, 08012, Spain*  
49  
50 43  
51 44 <sup>21</sup>*Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL 32611-0430,*  
52  
53 *USA*  
54  
55 45  
56 46 <sup>22</sup>*Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, College*  
57  
58  
59  
60

of Life Sciences, Beijing Normal University, Beijing, 100875, China

<sup>23</sup>State Key Laboratory of Biocontrol and College of Ecology and Evolution, Sun Yat-Sen University, Guangzhou, 510275, China

<sup>24</sup>Manomet Center for Conservation Sciences, Manomet, MA 02345, USA

<sup>25</sup>Schutzstation Wattenmeer Nationalparkhaus, Husum, D-25813, Germany

<sup>26</sup>Istituto Superiore per la Protezione e Ricerca Ambientale (ISPRA), Ozzano dell'Emilia BO, 40064, Italy

<sup>27</sup>School of Biology, University of St Andrews, St Andrews, KY16 9TH, United Kingdom

<sup>28</sup>Point Blue Conservation Science, Petaluma, CA 94954, USA

<sup>29</sup>Centre for Integrative Ecology, School of Life and Environmental Sciences, Faculty of Science, Engineering and the Built Environment, Deakin University, Burwood, 3125, Australia

<sup>30</sup>Quest University Canada, Squamish, 3200, Canada

<sup>31</sup>Department of Biology, University of Toliara, Toliara, Madagascar

**E-mail addresses:** orsolya.vincze@vocs.unideb.hu (OV), andras.kosztolanyi@gmail.com (AK),

barta.zoltan@science.unideb.hu (ZB), clemens.kuepper@uni-graz.at (CK), mm\_alrashedi@yahoo.com

(MA), amat@ebd.csic.es (JAA), araceli.a.tico@gmail.com (AAT), fionaburns\_21@hotmail.com (FB),

JCAVITT@weber.edu (JC), warren.conway@ttu.edu (WCC), tod472@gmail.com (MCL),

desucre@unam.mx (AEDM), ndosremedios@hotmail.co.uk (NdR), jordi@ebd.csic.es (JF),

dgalindoe@gmail.com (DGE), gegp01@gmail.com (GGP), salvachavin@hotmail.com (SGdA),

Cheri.Gratto-Trevor@ec.gc.ca (CGT), pauleric.jonsson@gmail.com (PJ), penn.lloyd@gmail.com (PL),

tmontal@aspb.cat (TM), jep33@bath.ac.uk (JEP), raya.pruner@gmail.com (RP), quepinjia@gmail.com

(PQ), liuy353@mail.sysu.edu.cn (YL), saalfeldst@gmail.com (STS), r.schulz@schutzstation-

wattenmeer.de (RS), lorenzo.serra@isprambiente.it (LS), jsc20@st-andrews.ac.uk (JSC),

1  
2 71 lstenzel@prbo.org (LS), mike.weston@deakin.edu.au (MAW), maiyasue@gmail.com (MY),  
3  
4 72 samazefania@yahoo.fr (SZ), T.Szekely@bath.ac.uk (TS)  
5  
6  
7 73  
8  
9 74 **Keywords**  
10 75 global change, climate, environmental stochasticity, seasonal environment, parental care, parental  
11 76 cooperation  
12  
13  
14 77  
15  
16  
17 78 **Running head:** Parental cooperation in fluctuating climate  
18  
19 79  
20  
21 80 **\*Correspondence to:** Orsolya Vincze, Email: orsolya.vincze@vocs.unideb.hu, Tel: +36-52-512900/  
22 81 62356  
23  
24 82 **Abstract word count:** 201  
25  
26 83 **Word count of the main body:** 5096  
27  
28  
29 84 **Number of references:** 56  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ABSTRACT

**Aim** Parental care improves offspring survival and therefore has a major impact on reproductive success. Whilst the influence of ambient environment on parental care is increasingly recognised, the impacts of environmental fluctuations remain largely unexplored. Assessing the impacts of environmental stochasticity, however, is essential for understanding how populations will respond to climate change. Here we investigate the influence of environmental stochasticity on biparental care in a worldwide avian genus.

**Location** Global

**Methods** We assembled data on biparental care in 36 plover populations (*Charadrius* spp.), from six continents, collected over several decades between 1981 and 2012. Using a space-for-time approach we investigate how average temperature, temperature stochasticity (i.e. year to year variation) and seasonality during the breeding season influences parental cooperation during care.

**Results** We show that both average ambient temperature and its fluctuations influence parental cooperation during incubation. Male care relative to females increases with both mean ambient temperature and stochasticity in temperature. Remarkably, local climatic conditions fully explained within-species, population differences in parental cooperation, but not differences among species.

**Main conclusions** Taken together, these results imply that climate change might have a multifaceted influence upon the reproductive behaviour and demography of populations by influencing parental care strategies and breeding systems.

## INTRODUCTION

Climate change influences the ecology and life-history of animals (Both & Visser 2001, Bradshaw & Holzapfel 2006; Dunn & Winkler 2010). It is associated with phenological shifts in life-history (e.g. earlier spring and/or later autumn migration, earlier breeding), changes in geographical ranges and



1  
2 109 physiology, as well as population trends (Walther *et al.*, 2002; Winkler *et al.*, 2002; Végvári *et al.*, 2010;  
3  
4 110 Thompson *et al.*, 2013; Vasseur *et al.*, 2014; IPCC, 2014; Lawson *et al.*, 2015). Although climate  
5  
6  
7 111 change has severe impacts on natural systems, our knowledge about how animals responds  
8  
9 112 behaviourally to altered climate is surprisingly limited. Monitoring behaviour would however enable us  
10  
11 113 to predict to what extent can behavioural plasticity mitigate the effects of climate change.  
12  
13  
14 114  
15  
16 115 Investigations of climate change often only focus on the impacts of average temperatures on  
17  
18 116 populations (Walther *et al.*, 2002). Nevertheless, there is a growing awareness that increased  
19  
20  
21 117 temperature variability, as well as a greater frequency and magnitude of climate extremes may also have  
22  
23 118 a significant effect on biological systems (Lawson *et al.*, 2015, Thompson *et al.*, 2013; Vasseur *et al.*,  
24  
25  
26 119 2014; IPCC 2014). Environmental uncertainty appears to increase with changing climatic conditions  
27  
28 120 (Thompson *et al.*, 2013; Vasseur *et al.*, 2014; Lawson *et al.*, 2015), therefore temperature fluctuations  
29  
30 121 may represent a potentially large, but to date mostly neglected threat to living organisms. In this study  
31  
32  
33 122 we aim to understand how animals respond to climate change in terms of behaviour and how  
34  
35 123 behavioural plasticity may mitigate the ecological impact of climate change. We investigate parental  
36  
37 124 care that is a major contributor to reproductive success in a wide range of taxa. Therefore, parental  
38  
39  
40 125 behaviour might represent an important link between climate change and its impacts on populations,  
41  
42 126 and it might change both in function of both average climatic conditions, as well as with its between-  
43  
44  
45 127 year and within-season variation (stochasticity and seasonality).  
46  
47 128  
48  
49 129 Parental care (i.e. parental behaviour that enhances the fitness of offspring and evolved for this  
50  
51  
52 130 function) is one of the most diverse social behaviours (Clutton-Brock 1991; McGraw *et al.*, 2010; Royle  
53  
54 131 *et al.*, 2012). There is immense variation in the type and duration of care parents provide, the timing and  
55  
56 132 duration of care-giving by each sex, and in ecological and morphological adaptations associated with  
57  
58  
59  
60

care (Clutton-Brock 1991; McGraw *et al.*, 2010; Royle *et al.*, 2012; Székely 2014; Bulla *et al.*, 2014).

Whilst parental behaviour has been studied extensively in wild populations (Royle *et al.*, 2012), evidence on how climate influences parental strategies is scant. Theoretical and empirical studies suggest that climate influences both the costs of care, i.e. the time and energy parents spend on rearing the young and the benefits of care, i.e. improved survival and recruitment of young (Clutton-Brock, 1991; Bonsall & Klug, 2011; Klug *et al.*, 2012). For instance, ambient temperature may influence the energetic costs of care (e.g. food provisioning, offspring brooding), and thus affect parental survival (Webb *et al.*, 2002; Bonsall & Klug, 2011; Klug *et al.*, 2012). Climatic conditions also influence the dependence of young on care, that particularly increases in extremely cold or hot climates, or during times of resource shortages. Parental protection and provisioning substantially improve offspring survival under these harsh conditions, as opposed to more favourable conditions (Wilson, 1975; Clutton-Brock, 1991; Alrashidi *et al.*, 2011, Bonsall & Klug, 2011). Although theoretical models suggest that increased climate variability will influence life-history trade-offs and thus parental care (Bonsall & Klug, 2011; Klug *et al.*, 2012; Tökölyi *et al.*, 2012), surprisingly little is known about the impact of these fluctuations on wild populations.

To explore the impact of climate on parental care, we investigate incubation behaviour, the most common form of care in birds (Deeming, 2002; Székely *et al.*, 2013). In nearly all bird species one (or both) parents incubate the eggs for several weeks, and in some cases for over two months (Deeming, 2002). By incubating the eggs, the parents keep egg temperature near the optimal for embryonic development by turning and warming or cooling the eggs in cold or hot climates, respectively (Deeming, 2002; AlRashidi *et al.*, 2011; Vincze *et al.*, 2013; Ghalambor & Martin, 2002; Royle *et al.*, 2012). Ambient temperature is expected to have a particularly significant impact on incubation in ground-nesting birds, because their eggs and the incubating parent are not buffered against extreme

1  
2 157 temperatures (Webb 1987; Deeming 2002; AlRashidi *et al.*, 2011).  
3  
4 158  
5  
6  
7 159 In environments with ambient temperatures close to optimal embryonic development (35-39°C: Webb,  
8  
9 160 1987), in the absence of other constraints, one parent may provide sufficient incubation (Deeming 2002;  
10  
11 161 AlRashidi *et al.*, 2011; Vincze *et al.*, 2013). If the environmental conditions, however, deviate from the  
12  
13  
14 162 optimal in either direction, one would expect increased parental effort by both sexes. However, male  
15  
16 163 involvement in parental care during incubation is usually less remarkable than that of females,  
17  
18 164 providing them with increased potential to alter their effort if needed (Auer *et al.*, 2007). Consequently,  
19  
20  
21 165 we expect males' share relative to females' to increase under harsh ambient conditions. Under harsh  
22  
23 166 environmental condition we mean high or low average temperatures or high interannual fluctuations of  
24  
25  
26 167 temperatures (stochasticity), since high between-year environmental fluctuations may increase the  
27  
28 168 probability of extreme events to occur. Additionally, we test the effect of within breeding season  
29  
30 169 environmental change. We predict increased male share in less seasonal, as well as in highly seasonal  
31  
32  
33 170 environments, in contrast to environments with medium seasonality. Under constant environmental  
34  
35 171 circumstances extended parental care is predicted for both sexes as part of the tropical life-history  
36  
37 172 syndrome (Wilson 1985), therefore in less seasonal environments male share should increase. Highly  
38  
39  
40 173 seasonal environments on the other hand restrict breeding time and remating opportunities, therefore it  
41  
42 174 might increase the value of current relative to future broods. Therefore, highly seasonal environments  
43  
44  
45 175 may also select for increased male share relative to females.  
46  
47 176  
48  
49 177 In this study we use data from 36 plover populations. Plovers (*Charadrius spp.*) are ground nesting  
50  
51  
52 178 shorebirds with body mass ranging from approximately 20g to 50g. The ancestor of this monophyletic  
53  
54 179 group likely evolved in temperate or cold climates of the Northern hemisphere (dos Remedios *et al.*  
55  
56 180 2015). Plovers breed on all continents except Antarctica in habitats as varied as arctic tundra, temperate  
57  
58  
59  
60

grassland, tropical beaches, salt marshes, sand dunes, semi-deserts, deserts and high altitude mountain lake shores (Piersma & Wiersma 1996). This immense variation in breeding environment provides an excellent opportunity to conduct a geographically large-scale study, capturing a substantial range of global ecological diversity. Plovers usually lay 2-4 eggs in uninsulated scrapes. Incubation is usually carried out by both parents, although the extent of male involvement in incubation is highly variable among species and populations (Vincze *et al.*, 2013). In addition, the share of incubation by each sex may vary throughout the day: in most species males tend to incubate at night, whereas females carry out most of the daytime incubation (Vincze *et al.*, 2013; but see St Clair *et al.*, 2010a).

Here we investigate how climate influences parental behaviour using an extensive data set on parental care that cover temperate and tropical habitats in both the northern and southern hemispheres (between 55°N to 52°S latitude, and between 145°E to 121°W longitude). To see how climate influence incubation behaviour, we used the space-for-time substitution approach, i.e. we infer temporal trends from spatial data, a powerful method in ecology (Pickett, 1989). First, we establish that how the division of parental care varies across species, populations and over the day. Second, we test whether ambient temperature and fluctuations in temperature influence the division of care between males and females.

## METHODS

### Fieldwork

Fieldwork was carried out in 36 breeding populations, and ranged from one to 16 breeding seasons per population (Table S1). Parents were captured on their nest using funnel traps, noose mats, box traps or bownet traps while incubating (see Székely *et al.*, 2008 for general methodology, and specific references in Table S1). For each captured bird we recorded the time of capture and sex of the captured individual. In three populations (Florida, Monterey Bay, Cape Peninsula) capture data were augmented

1  
2 205 by opportunistic observations of the incubating parent. Sex determination was based on plumage  
3  
4 206 characteristics in the field and/or measurements (e.g. vent), sex-specific DNA markers (following  
5  
6 207 methods in Parra *et al.*, 2014; Gratto-Trevor, 2011), and, in a few cases, based on observations of  
7  
8 208 copulation behaviour (Table S1).  
9  
10  
11 209  
12  
13  
14 210 Egg-laying date was defined as the date of clutch completion. This was either known, for nests that  
15  
16 211 were found during egg-laying, or estimated by floating eggs or measuring egg mass relative to egg size  
17  
18 212 (Székely *et al.*, 2008; Fraga & Amat, 1996). Egg-laying dates were standardised separately for each  
19  
20 213 population by subtracting the mean and dividing by the standard deviation of laying dates for a given  
21  
22 214 population. Since males have a greater tendency to be at the nest during egg-laying and egg-hatching  
23  
24 215 (Székely T & Kosztolányi A, pers. obs), we only included nests that were incubated for at least three  
25  
26 216 days and but not longer than 20 days (incubation usually lasts for 25-26 days in small plovers, Piersma  
27  
28 217 & Wiersma, 1996). If an individual was captured (or observed) several times, we only included its first  
29  
30 218 record, in order to exclude birds with potentially altered behaviour due to previous disturbance. To  
31  
32 219 investigate daily patterns of incubation behaviour, we divided the day into twelve 2-hour time periods  
33  
34 220 following previous analyses of incubation patterns in small plovers (AlRashidi *et al.*, 2011; Vincze *et*  
35  
36 221 *al.*, 2013). Records between 00h and 04h were not included in data analyses, since we lacked such data  
37  
38 222 from most populations. To estimate parental care division between the sexes, we used the sex of  
39  
40 223 incubating parent as binary response variable in statistical models. In total, 5,591 individuals were  
41  
42 224 included in the dataset (Table S1).  
43  
44  
45  
46  
47  
48  
49 225  
50

51  
52 226 **Consistency between captures and behavioural observations**  
53

54 227 To test whether capture times reflected the daily routine of shared incubation between the sexes, we  
55  
56 228 compared male share as estimated based on capture data with male share as estimated based on  
57  
58  
59  
60

continuous behavioural observations in six populations of two species, from which both capture data and behavioural data were available (see Vincze *et al.*, 2013 for details on behavioural observations). Based on capture data, male share (% capture) was calculated as the percentage of male captures of all captures (males plus females) at the nests during a given 2-hour time period. Based on behavioural observations, male share (% behaviour) was calculated as the % of time when males incubated of the total time the nest was incubated by either parent in a given 2-hour time period. The relationship between capture-based and behavioural observation-based male share estimates was analysed using linear regressions for the six populations separately, where each 2-hour time period represented a datum. These data points were weighted by the number of captures in each 2-hour time period, since the precision of the male share (% capture) estimate is expected to increase with the total number of individuals captured in a given time period.

#### Climate data

We extracted ambient temperature data from the University of East Anglia Climate Research Unit database (CRU, <http://www.cru.uea.ac.uk/>; version 3.10.01; Mitchell & Jones, 2005). The CRU database is a global dataset containing interpolated monthly average temperatures (°C) from 1901 onward in a grid of spatial coordinates (0.5 x 0.5 degrees). For each population we selected temperatures from 20 years prior to the last year of data collection (including the latter); this seemed sufficient to represent the ambient temperatures the plovers experienced in our study given that the largest temporal dataset based on captures spanned 16 years. Since our study focused on parental behaviour, we only used ambient temperatures from those months when capture data were collected in each population; these months are referred to as 'the breeding season'. Using the same number of years for each population enabled us to estimate the three climate variables used here (see below) with similar precision in each population, irrespective of the number of data collection years in each of these.

1  
2 253 Note that although results presented are based on climate data of 20 years, we carried out sensitivity  
3  
4 254 analyses by repeating the analyses using 15, 10 and 5 years climate data prior to the last year of field  
5  
6  
7 255 data collection. These models yielded highly consistent results (see SI Appendix, Table S2).  
8  
9 256  
10  
11 257 We derived three variables to characterise ambient environment. (i) Average temperature at each site  
12  
13 258 refers to mean temperature over the breeding season, calculated from monthly means for each breeding  
14  
15  
16 259 season and averaged over 20 years. (ii) Between-year variation was calculated as the standard deviation  
17  
18  
19 260 of each month's average temperature across the 20 years, averaged over the breeding season for each  
20  
21 261 population. (iii) Within-season temperature variation was obtained by calculating the average  
22  
23 262 temperature for each month of the breeding season over 20 years, and then calculating the difference  
24  
25  
26 263 between the maximum and minimum monthly average temperatures. Therefore, the latter two variables  
27  
28 264 refer to the average between-year and within-season variation in ambient temperature during breeding at  
29  
30  
31 265 a given site. Climate variables tend to be correlated (see for example Tökölyi *et al.*, 2014), therefore to  
32  
33 266 test whether collinearity exists in models containing all three temperature variables, we calculated  
34  
35 267 variance inflation factors (VIFs) for models without interactions, using the "vif.mer" function (available  
36  
37 268 at: <https://github.com/aufrank/R-hacks/blob/master/mer-utils.R>, last accessed on: 15 September 2014) in  
38  
39  
40 269 R (R Core Team, 2014). None of the three climatic variables had  $VIF > 2.52$ . Additionally, none of the  
41  
42 270 correlation coefficients between pairs of climate variables across populations exceeded 0.55 (Pearson  
43  
44  
45 271 correlation). Therefore, collinearity between temperature variables does not seem to be a major issue in  
46  
47 272 our analyses.  
48  
49 273

50  
51  
52 274 **Statistical analyses**  
53

54 275 Since no population-level phylogenetic hypothesis is available for the 36 plover populations studied  
55  
56 276 here, we used mixed-effects models to analyse relationships between care division and environmental  
57  
58  
59  
60



data. To account for the phylogenetic non-independence we included population and species identity as random factors. We used the sex of parents (male or female) captured on the nest as the response variable in binomial models. Species, population and nest identity were included as nested random factors in all models. Although we only used one capture per individual, nest identity was included as a random factor in the models to control for potential non-independence of male and female behaviour for a given nest. Time period was included in models as a fixed factor with 10 levels (i. e. 2 hour windows, between 04h and 24h). The three temperature variables were standardised, using the “scale” function implemented in R, to ease model fitting and comparing the effects. The standardised variables were included in the models as second order orthogonal polynomials, because of the expected non-linear effects (see above, Vincze *et al.* 2013). Although we also tested the effects of laying date, it was excluded from further models because it did not correlate with the sex of the parent captured.

We built four mixed effects models. First, to test how care division varies throughout the day and across species and populations we constructed a model that included time period and the random factors of species, population and nest IDs (model 1). Second, to test whether the daily pattern of incubation differed between plover species and populations, we built two models: in one of these models additionally to the terms in model 1 we included the species x time period interaction (model 2), whereas in the other the population x time period interaction was included (model 3).

Third, to investigate the effects of ambient temperature, and its fluctuations between years and within-seasons, we constructed a model (model 4) that included the time period factor, the three temperature variables (i.e., mean, within-season and between year variation), and two-way interactions between the time period and temperature variables. The significance of each predictor was assessed by removing it from the model and comparing the resulting model to the original using likelihood ratio statistics (see



1  
2 301 Figure S1 for schematic illustration of hypothesis testing).  
3  
4 302  
5  
6  
7 303 Mixed models were built using the “glmer” function, as implemented in the “lme4” package (version  
8  
9 304 1.1-7, Bates *et al.*, 2015) in R (version 3.1.1, R Core Team, 2014).  
10  
11 305

12  
13  
14 306 **Daily routines of parental care in different climate scenarios**

15  
16 307 To investigate the impact of climate on daily routines during incubation, we removed from model 4 the  
17  
18 308 non-significant interaction and quadratic terms for between-year variation (Table 1), and used this  
19  
20 309 resulting model for predictions. We predicted the effect of the three temperature variables on daily  
21  
22 310 routines of care division for nine climate scenarios. For each temperature variable, we calculated the  
23  
24 311 predicted values for the 10 time periods at the 2.5% quantile, median and 97.5% quantile value of the  
25  
26 312 temperature variable in question, while the other two temperature variables were kept at their median  
27  
28 313 values. Only fixed effects were taken into account when extracting model predictions.  
29  
30  
31  
32  
33 314

34  
35 315 **RESULTS**

36  
37 316 **Consistency between captures and behavioural observations**

38  
39 317 Capture-based behavioural estimates reflect parental care division in plovers, since capture-based  
40  
41 318 estimates of male share were highly correlated with estimates of male share obtained by behavioural  
42  
43 319 observations (Figure 1,  $R^2 = 0.61 - 0.97$ ,  $n = 6$  populations).  
44  
45  
46  
47 320

48  
49 321 **Incubation routines in different populations**

50  
51 322 Incubation routines differed between different plover species and populations (models 2 & 3, Table 1).  
52  
53 323 On the one hand, in species like *C. melodus*, males and females spent comparable time on incubation  
54  
55 324 throughout the day (Figure 2). On the other hand, incubation routines followed a diurnal pattern in  
56  
57  
58  
59  
60

species such as *C. alexandrinus*, *ruficapillus* and *modestus* (Figure 2). Furthermore, there were considerable differences in daily pattern of incubation among the different populations of the same species (Figure 2).

### **Ambient environment, between- and within-season variation**

Mean ambient temperature, as well as between- and within-season variation in temperature strongly influenced parental care division (model 4, Table 1). Male share of incubation generally increased with mean ambient temperature. This effect was, however, dependent on time of the day as indicated by the significant interaction between time period and mean ambient temperature. For example, during daylight hours (8 - 20 h) males' share of incubation increased with mean ambient temperature, though the increase was non-linear and varied depending on the specific time window (Figure 3a).

Temperature fluctuations also predicted incubation (Figure 3b and c). Between-year variation tended to have a linear influence on daily shifts: male share of incubation increased with variation in temperature between years and this effect was similar throughout the day (Figure 3b). Within-season temperature change also predicted shifts in daily routines of males relative to females: with increasing change in temperature during the breeding season, male share decreased between 6 h and 16 h. The effect of within-season temperature variation was however strongly non-linear early in the morning and in the evening (Figure 3c).

Once the three temperature variables were included in the models, the variance explained by population decreased considerably from 0.115 (model 1) to 0.005 (model 4). In contrast, the variance explained by species changed very little from 0.184 (model 1) to 0.191 (model 4).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Daily routines in different climate scenarios**

With increasing mean ambient temperature and between-year variation, male share increases during daylight hours, while in the case of mean temperate this happened at the expense of a lowered share of care during the early morning hours (Figure 4a,b). Furthermore, with increasing within-season temperature variation, male share in incubation decreases during daylight hours (Figure 4c).

**DISCUSSION**

Three major insights have emerged from our study regarding the effect of environment on parental behaviour. First, male contribution to parental care was strongly influenced by ambient temperature. Second, temperature effects on behaviour varied with time of the day. Therefore, not just overall care division changed with changing environmental conditions, but the daily routine of care division was also affected. Specifically, male share of parental care increased with mean temperature and between-year variation in temperature during daylight hours. When conditions became harsher, i.e. the mean temperature and or the between-year unpredictability of temperature was high, males generally increased their effort relative to females during incubation. Finally, geographic variation in care division within species was largely explained by local ambient temperatures, although the differences between different species persisted even after controlling for climatic effects. The latter suggests that different plover populations respond in similar ways to ambient environment, reflecting phenotypic plasticity in behaviour. In contrast, there is substantial species difference in parental care, reflecting a strong phylogenetic effect.

Our results highlight that not only the average environmental conditions, but also their between- and within-season variation play a pivotal role in shaping care division and daily routines of parental care in biparental species. Environmental uncertainty influences reproduction (e.g. breeding initiation, song

display) and life-history (e.g. egg size, clutch size, age of sexual maturity; Lips, 2001; Dewar & Richard, 2007; Botero *et al.*, 2009; Bonsall & Klug, 2011). In addition, unpredictable environmental variation influences mating systems (Botero & Rubenstein, 2012), and may promote the evolution of cooperative breeding strategies (Rubenstein & Lovette, 2007; Jetz & Rubenstein, 2011; but see Gonzalez *et al.*, 2013 for opposite effect). Here we show that parental cooperation is also strongly influenced by predictable and stochastic climate variations.

We propose that more cooperative male behaviour is driven by the need to protect the embryo better under higher frequencies of extreme events (Deeming, 2002; AlRashidi *et al.*, 2011). The expected changes in care division are most likely to occur during mid-day leading to altered daily routines of parental care. As climate change models predict both an increase in temperature and greater frequency of extreme events (Vasseur *et al.*, 2014; IPCC, 2014; Lawson *et al.*, 2015), our findings suggest that pattern of parental care will shift in the near future in biparental species. Such shifts may include greater diurnal incubation responsibilities for the sex with the more variable parental contribution (usually males in birds and mammals, Clutton-Brock, 1991). On the one hand, these shifts may help to maintain hatching success and hatchling condition in the provisioned brood under worsening environmental conditions (Reid *et al.*, 2002). On the other hand, they may preclude the sex that increases parental effort from performing other activities (Dunning, 2002; Reid *et al.*, 2002; Bulla *et al.*, 2014). For instance, a greater share of care division by a given sex may constrain its foraging time, or may reduce its ability to attract further mates or provision other broods, therefore may directly influence mating systems (e.g. Reid *et al.*, 2002). These effects would be especially important in species with flexible and variable parental care and mating systems (e.g. Reid *et al.*, 2002; Kosztolányi *et al.*, 2006).

Periodicity over the day drives daily behavioural routines (Houston & McNamara 1999). Similar to

1  
2 397 earlier studies (AlRashidi *et al.* 2011, Vincze *et al.*, 2013), we found significant daily variation in care  
3  
4 398 provisioning by each sex in specific plover populations. A novel aspect of our current study is that we  
5  
6  
7 399 relate variations in daily routines of care to variation in environmental variables. Our results suggest  
8  
9 400 that behavioural response to temperature changed during the day, in particular, behaviour around mid-  
10  
11 401 day seemed to be most influenced. This suggests that breeding routines are driven by avoiding extreme  
12  
13  
14 402 hot temperatures. These results may contribute to a detailed theoretical treatment of daily parental  
15  
16 403 routines. The current lack of such models hampers our ability to provide a more detailed explanation for  
17  
18  
19 404 the effect of environmental conditions on daily routines and hence to guide further empirical  
20  
21 405 investigations.  
22  
23 406  
24  
25 407 Since male contribution to care correlates with other aspects of breeding systems (e.g. 0% male care  
26  
27  
28 408 usually associated with polygyny, whereas 100% male care may be associated with polyandry and sex  
29  
30  
31 409 role reversal, Searcy & Yasukawa, 1995, Liker *et al.*, 2013), our work suggests that breeding systems  
32  
33 410 will also respond to changes in ambient temperature. To follow up this line of investigation, it would be  
34  
35 411 interesting to study how brood care patterns, frequency of polygamy and extra pair paternity may vary  
36  
37  
38 412 in relation to environmental fluctuations (e.g. in temperatures, food, resource quality, and territory  
39  
40 413 quality). Since these reproductive behaviours make fundamental contributions to reproductive success,  
41  
42 414 we believe it is imperative to assess the impact of climate change not only on parental behaviour, but on  
43  
44  
45 415 other aspects of breeding systems including mate choice, mating system, and pair bonding.  
46  
47 416  
48  
49 417 Care division within a species varied with between-population differences in climatic conditions. Local  
50  
51  
52 418 adaptation is unlikely since many plover species show low genetic differentiation (Funk *et al.* 2007,  
53  
54 419 Küpper *et al.*, 2012, Eberhardt-Phillips *et al.*, 2015). It is more likely that sex roles during biparental  
55  
56  
57 420 care are phenotypically plastic within species, and are modulated by local conditions. This interpretation  
58  
59  
60

is consistent with previous studies, which have demonstrated behavioural plasticity according to local environments during incubation (Al Rashidi *et al.*, 2011, Vincze *et al.*, 2013). Another consequence of the observed flexibility in parental behaviour is that these populations might effectively be able to cope with changing climate at least within the climate range studied here. More climate resilience may be achieved by phenological changes (e.g. Chambers *et al.*, 2008).

Although we found highly significant relationships between environmental fluctuation, its within-year variability, parental care division and its daily routines, the theoretical bases of these relationships have not been fully explored (Klug *et al.*, 2012). Previous theoretical analyses of care and life history traits pointed out that environmental unpredictability can have sophisticated and counter-intuitive influences on provision of care (Klug *et al.*, 2012). To model these future scenarios, it is essential to assess how different aspects of climate influence contemporary populations. Since changing climate may alter the costs and benefits of parental care (Clutton-Brock, 1991; Royle *et al.*, 2012; IPCC, 2014), climate change is likely to affect the reproductive success of individuals that, in turn, will be likely to have an impact on population growth and resilience. We call for new theoretical models to tease apart the effects of ambient environment, social environment and life-histories on care provisioning and its daily pattern.

Using parental care data from an exceptionally wide geographic range, we have shown that incubation, a major component of parental care in birds, is significantly related to mean and variation of ambient temperatures. Daily patterns of care division between the sexes are strongly affected by temperatures. Theoretical explorations show that ambient temperature, as well as its predictable and unpredictable fluctuations, will influence diurnal incubation patterns (Bonsall & Klug, 2011; Klug *et al.*, 2012). We recommend follow up studies building upon our research framework by augmenting these analyses with other climatic variables (e.g. precipitation, wind), and using a variety of response variables such as

1  
2 445 mating system, brood survival and life-histories. In addition, we encourage the development of  
3  
4 446 theoretical models investigating the influence of environmental fluctuations on parental care and  
5  
6  
7 447 breeding system.  
8  
9 448  
10  
11 449 **ACKNOWLEDGEMENTS**  
12  
13  
14 450 Fieldwork and blood sampling was authorized by relevant authorities: Hungary (Environmental  
15  
16 451 Ministry and Kiskunság National Park), Australia (Department of Primary Industries and Environment  
17  
18 452 and Parks Victoria, #F10005850), Argentina (Centro Nacional Patagonico, Dr. Luis Bala), Ceuta  
19  
20  
21 453 (#SGPA/DGVS/01367/11), San Quintín Bay (Dirección General de Vida Silvestre  
22  
23 454 SGPA/DGVS/02078/12), Llobregat Delta (Ministry of Environment #660117), Fuente de Piedra  
24  
25  
26 455 (Consejería de Medio Ambiente, Junta de Andalucía), Farasan (Saudi Wildlife Commission), California  
27  
28 456 (U.S. Fish and Wildlife (USFWS) #TE807078 and U. S. Geological Survey (USGS) #09316),  
29  
30 457 Madagascar (Ministry of Environment, Forests and Tourism of the Republic of Madagascar  
31  
32  
33 458 #053/11/MEF/SG/DGF/DCB.SAP/SCB and #132/10/MEF/SG/DGF/DCB.SAP/SSE; Service de la  
34  
35 459 Gestion de la Faune et de la Flore, Direction de la Valorisation des Ressources Naturelles, Ministère de  
36  
37 460 l'Environnement et des Forêts Madagascar #080N-EA06/MG11; Madagascar National Parks #398-  
38  
39  
40 461 10/MEF/SG/DGF/DVRN/ SGFF), Falkland Islands (Falkland Islands Government, #R08/2007),  
41  
42 462 Canada (Environment Canada, Agriculture Canada), China (Hebei Forestry Bureau). Souh-Africa (Cape  
43  
44  
45 463 Nature and SAFRING), Nayarit (#USFWS 23021), Maio (Directorate Geral Ambiente), Tuzla (Turkish  
46  
47 464 Ministry of National Parks, Tuzla Municipality and Governor of Karatas, Mr. E. Karakaya), Germany  
48  
49 465 (Institut für Vogelforschung, "Vogelwarte Helgoland", Wilhelmshaven), Florida (Institutional Animal  
50  
51  
52 466 Care and Use Committee at the University of Florida, project no. E877) and all banding was completed  
53  
54 467 under banding permit #21980 in collaboration with the Florida Fish and Wildlife Conservation  
55  
56 468 Commission, Texcoco (SEMARNAT/SGPA/DGVS/00098/14), Texas (Institutional Animal Care and  
57  
58  
59  
60



Use Committee at Stephen F. Austin State University #TECMW 10-08-07; U.S. Geological Survey Master Banding Permit #23393; Texas Parks and Wildlife Department SPR-0804-009), St Helena (St Helena Government and the BTO, permit number C5326). OV was supported by the European Union and the State of Hungary, co-financed by the European Social Fund in the framework of TÁMOP-4.2.4.A/ 2-11/1-2012-0001 'National Excellence Program' and by the Hungarian Ministry of Human Resources, National Talent Program (NTP-EFÖ-P-15-0043) during the writing of the manuscript. AK was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. MAW was funded by the Hermon Slade Foundation, BirdLife Australia, M.A. Ingram Trust, and prominent volunteer input from Laura Tan, Kasun Ekanayake, Stephanie Lomas, and Jessica Bywater. MY was funded by NSERC PhD fellowships and a SSHRC research grant. CK was funded by a Marie Curie IEF postdoctoral Fellowship. MCL, DG, GEGP were funded by CONACyT graduate programmes, PQ was funded by United Foundation for Natural Science of National Natural Science Foundation of China and People's Government of Guangdong Province (No. U0833005) and YL was funded by the open project of State Key Laboratory of Biocontrol, Sun Yat-sen University. We thank Drs. Luis Bala and María de los Ángeles Hernández, for their support and advice during fieldwork in Argentina. Details about support for the study at Ceuta Bay is listed on the website [chorlito.org](http://chorlito.org). Madagascar plover fieldwork and molecular sexing were supported by Leverhulme Trust (ID200660763, Ecology and genetics of endemism) and NERC Biomolecular Analysis Facility at the University of Sheffield respectively. JSC was supported by the University of Bath and Falkland Islands Government, and assisted in the field by Neil Dawson, Chris Dodd, Philipp Herrmann and Jacqueline Pearson. JAA was supported by research projects PB95-0110 and CGL2011-24230 from Ministerio de Ciencia e Innovación, Spain, both with EU-ERDF financial support. DGE was funded by the Sonoran Joint Venture (FWS Agreement No. F11AP00052), and by graduate scholarships from the PIFI at Instituto Politécnico Nacional-CICIMAR (20120989) and the Mexican National Council for Science and Technology (CONACYT, No. 99866).



1  
2 493 FB was funded by the University of Bath and the RSPB, with support from St Helena National Trust.  
3  
4 494 CLGT was funded by Environment Canada (Species at Risk), and the Government of Canada's  
5  
6 495 Interdepartmental Recovery Fund. ZB was supported by a NKFIH grant (grant no. K112527). Financial  
7  
8  
9 496 support: Florida (US Fish and Wildlife Service Panama City Field Office), Germany (Stifterverband für  
10  
11 497 die Deutsche Wissenschaft, Essen), Nayarit (Tides Foundation, Rio Tinto Kennecott, BirdLife  
12  
13  
14 498 International, US Fish and Wildlife Service Region 6, Denker's Family Foundation and Weber State  
15  
16 499 University). This study is contribution 12319 of Point Blue Conservation Science.  
17  
18  
19 500

20  
21 501 **SUPPORTING INFORMATION**

22  
23 502 **Figure S1** Schematic illustration of hypothesis testing.

24  
25 503 **Table S1** Summary of the data used.

26 504 **Table S2** Sensitivity analyses.

27  
28 505 **Appendix S1** Supporting references.  
29  
30  
31  
32  
33 506

34  
35 507 **BIOSKETCH**

36  
37 508 Orsolya Vincze is a PhD candidate at the University of Debrecen. Her research focuses on behavioural  
38  
39 509 ecology and ecophysiology of birds.  
40  
41  
42 510

43  
44 511 **REFERENCES**

45  
46 512 AlRashidi, M., Kosztolányi, A., Shobrak, M., Küpper, C. & Székely T. (2011) Parental cooperation in  
47  
48 513 an extreme hot environment: natural behaviour and experimental evidence. *Animal Behaviour*, **82**,  
49  
50  
51 514 235–243.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- Auer, S.K., Bassar, R.D. & Martin, T.E. (2007) Biparental incubation in the chestnut-backed tit-babbler *Parisoma subcaeruleum*: mates devote equal time, but males keep eggs warmer. *Journal of Avian Biology*, **38**, 278-283.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1-48.
- Bonsall, M.B. & Klug, H. (2011) The evolution of parental care in stochastic environments. *Journal of Evolutionary Biology*, **24**, 645-655.
- Both, C. & Visser, M.E. (2001) Adjustment to climate change is constrained by arrival date in a long-distance migrant bird. *Nature*, **411**, 296-298.
- Botero, C.A., Boogert, N.J., Vehrencamp, S.L. & Lovette, I.J. (2009) Climatic patterns predict the elaboration of song displays in mockingbirds. *Current Biology*, **19**, 1151-1155.
- Botero, C.A. & Rubenstein, D.R. (2012) Fluctuating environments, sexual selection and the evolution of flexible mate choice in birds. *PLoS ONE*, **7**, e32311.
- Bradshaw, W.E. & Holzapfel, C.M. (2006) Evolutionary response to rapid climate change. *Science*, **312**, 1477-1478.
- Bulla, M., Valcu, M., Rutten, A.L. & Kempenaers, B. (2014) Biparental incubation patterns in a high-Arctic breeding shorebird: how do pairs divide their duties? *Behavioral Ecology*, **25**, 152-164.
- Chambers, L.E., Gibbs, H., Weston, M.A. & Ehmke, G.C. (2008) Spatial and temporal variation in the breeding of masked lapwings (*Vanellus miles*) in Australia. *Emu*, **108**, 115-124.
- Clutton-Brock, T.H. (1991) *The evolution of parental care*. Princeton University Press, Princeton.
- Deeming, C. (2002) *Avian incubation: behaviour, environment and evolution*. Oxford University Press, Oxford.

1  
2 537 Dewar, R.E. & Richard, A.F. (2007) Evolution in the hypervariable environment of Madagascar.  
3  
4 538 *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 13723-  
5  
6 539 13727.  
7  
8  
9 540 Dickey, M.H., Gauthier, G. & Cadieux, M.C. (2008) Climatic effects on the breeding phenology and  
10  
11 541 reproductive success of an arctic nesting goose species. *Global Change Biology*, **14**, 1973-1985.  
12  
13  
14 542 Dos Remedios, N., Lee, P.L., Burke, T., Székely, T. & Küpper, C. (2015) North or south? Phylogenetic  
15  
16 543 and biogeographic origins of a globally distributed avian clade. *Molecular Phylogenetics and*  
17  
18 544 *Evolution*, **89**, 151-159.  
19  
20  
21 545 Dunn, P.O. & Winkler, D.W. (2010) Effects of climate change on timing of breeding and reproductive  
22  
23 546 success in birds. In: *Effects of climate change on birds* (eds. Møller, A.P., Fiedler, W. & Berthold,  
24  
25 547 P.), PP. 113-128. Oxford University Press, Oxford.  
26  
27  
28 548 Eberhart-Phillips, L.J., Hoffman, J.I., Brede, E.G., Zefania, S., Kamrad, M., Székely, T. & Bruford,  
29  
30 549 M.W. (2015) Contrasting genetic diversity and population structure among three sympatric  
31  
32 550 Madagascan shorebirds: parallels with rarity, endemism, and dispersal propensity. *Ecology and*  
33  
34 551 *Evolution*, **5**, 997-1010.  
35  
36  
37 552 Fraga, R.M. & Amat, J.A. (1996) Breeding biology of a Kentish plover (*Charadrius alexandrinus*)  
38  
39 553 population in an inland saline lake. *Ardeola*, **43**, 69-85.  
40  
41  
42 554 Funk, W.C., Mullins, T.D. & Haig, S.M. (2007). Conservation genetics of snowy plovers (*Charadrius*  
43  
44 555 *alexandrinus*) in the Western Hemisphere: population genetic structure and delineation of  
45  
46 556 subspecies. *Conservation Genetics*, **8**, 1287-1309.  
47  
48  
49 557 Ghalambor, C.K. & Martin, T.E. (2002) Comparative manipulation of predation risk in incubating birds  
50  
51 558 reveals variability in the plasticity of responses. *Behavioral Ecology*, **13**, 101-108.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- Gonzalez, J.C.T., Sheldon, B.C. & Tobias, J.A. (2013) Environmental stability and the evolution of cooperative breeding in hornbills. *Proceedings of the Royal Society of London Series B-Biological Sciences*, **280**, 20131297.
- Gratto-Trevor, C.L. (2011) Ageing and sexing the Piping Plover *Charadrius melodus*. *Wader Study Group Bulletin*, **118**, 118–122.
- Houston, A.I. & McNamara, J.M. (1999) *Models of adaptive behaviour: an approach based on state*. Cambridge University Press, Cambridge.
- IPCC (2014) Climate change 2014: impact adaptation and vulnerability. AR5, Working Group II (WGII AR5) Available at: <http://ipcc-wg2.gov/AR5/>. Last accessed 10 July 2014.
- Jetz, W. & Rubenstein, D.R. (2011) Environmental uncertainty and the global biogeography of cooperative breeding in birds. *Current Biology*, **21**, 72–78.
- Klug, H., Alonzo, S.H. & Bonsall, M.B. (2012) Theoretical foundations of parental care. In: *The evolution of parental care* (eds. Royle, N.J., Smiseth, P.T. & Kölliker, M.), PP. 21–39. Oxford University Press, Oxford.
- Kosztolányi, A., Székely, T., Cuthill, I.C., Yilmaz, K.T. & Berberoglu, S. (2006) Ecological constraints on breeding system evolution: the influence of habitat on brood desertion in Kentish plover. *Journal of Animal Ecology*, **75**, 257–265.
- Küpper, C., Edwards, S.V., Kosztolányi, A. *et al.* (2012). High female mediated gene flow 1 on a continental scale in the polyandrous Kentish Plover *Charadrius alexandrinus*. *Molecular Ecology*, **21**, 5864–5879.
- Lawson, C.R., Vindenes, Y., Bailey, L. & Pol, M. (2015) Environmental variation and population responses to global change. *Ecology letters*, **7**, 724–736.
- Liker, A., Freckleton, R.P. & Székely, T. (2013) The evolution of sex roles in birds is related to adult sex ratio. *Nature Communications*, **4**, 1587.

1  
2 583 McGraw, L., Székely, T. & Young, L.J. (2010) Pair bonds and parental behaviour. In: *Social behaviour:*  
3  
4 584 *genes, ecology and evolution* (eds. Székely, T., Moore, A. & Komdeur, J.), PP. 271–301,  
5  
6 585 Cambridge University Press, Cambridge.  
7  
8  
9 586 McNamara, J.M., Mace, R. & Houston, A.I. (1987) Optimality daily routines of singing and foraging in  
10  
11 587 a bird singing to attract a mate. *Behavioral Ecology and Sociobiology*, **20**, 399-405.  
12  
13  
14 588 Mitchell, T.D. & Jones, P.D. (2005) An improved method of constructing a database of monthly climate  
15  
16 589 observations and associated high-resolution grids. *International journal of climatology*, **25**, 693-  
17  
18 590 712.  
19  
20  
21 591 Pickett, S.T.A. (1989) Space-for-time substitution as an alternative to long-term studies. In: *Long-Term*  
22  
23 592 *Studies in Ecology: Approaches and Alternatives* (ed. Likens, G.E.), PP. 110-135. Springer, New  
24  
25 593 York.  
26  
27  
28 594 Piersma, T. & Wiersma, P. (1996) Family *Charadriidae* (plovers). In: *Handbook of the Birds of the*  
29  
30 595 *World*, Vol. 3, (eds. del Hoyo, J., Elliott, A. & Sargatal, J.), PP. 384-442. Lynx Editions,  
31  
32 596 Barcelona.  
33  
34  
35 597 Parra, J.E., Beltrán, M., Zefania, S., Dos Remedios, N. & Székely, T. (2014) Experimental assessment  
36  
37 598 of mating opportunities in three shorebird species. *Animal Behaviour*, **90**, 83-90.  
38  
39  
40 599 R Core Team (2014) R: A language and environment for statistical computing. R Foundation for  
41  
42 600 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.  
43  
44  
45 601 Reid, J.M., Monaghan, P. & Ruxton, G.D. (2002) Males matter: the occurrence and consequences of  
46  
47 602 male incubation in starlings (*Sturnus vulgaris*). *Behavioral Ecology and Sociobiology*, **51**, 255-  
48  
49 603 261.  
50  
51  
52 604 Royle, N.J., Smiseth, P.T. & Kölliker, M. (2012) *The evolution of parental care*. Oxford University  
53  
54 605 Press, Oxford.  
55  
56  
57  
58  
59  
60

- Rubenstein, D.R. & Lovette, I.J. (2007) Temporal environmental variability drives the evolution of cooperative breeding in birds. *Current Biology*, **17**, 1414-1419.
- Searcy, W.E. & Yasukawa, K. (1995) Polygyny and sexual selection in red-winged blackbirds. Princeton University Press, Princeton, NJ.
- St Clair, J.J.H., Küpper, C., Herrmann, P., Woods, R.W. & Székely, T. (2010a). Unusual incubation sex-roles in the Rufous-chested Dotterel *Charadrius modestus*. *Ibis*, **152**, 402.
- Székely, T., Kosztolányi, A. & Küpper, C. (2008) Practical guide for investigating breeding ecology of Kentish plovers *Charadrius alexandrinus*. Version 3. Unpublished report, University of Bath.
- Available at: [http://www.bath.ac.uk/bio-sci/biodiversity-lab/pdfs/KP\\_Field\\_Guide\\_v3.pdf](http://www.bath.ac.uk/bio-sci/biodiversity-lab/pdfs/KP_Field_Guide_v3.pdf). Last accessed 23 July 2014.
- Székely, T., Remeš, V., Freckleton, R.P. & Liker, A. (2013) Why care? Inferring the evolution of complex social behaviour. *Journal of Evolutionary Biology*, **26**, 1381-1391.
- Székely, T. (2014) Sexual conflict between parents: offspring desertion and asymmetrical parental care. In: *Sexual conflict* (eds. Gavrillets, S & Rice, W.), Cold Spring Harbor Perspectives in Biology.
- Thompson, R.M., Beardall, J., Beringer, J., Grace, M. & Sardina, P. (2013) Means and extremes: building variability into community-level climate change experiments. *Ecology Letters*, **16**, 799-806.
- Tökölyi, J., McNamara, J.M., Houston, A.I. & Barta, Z. (2012) Timing of avian reproduction in unpredictable environments. *Evolutionary Ecology*, **26**, 25-42.
- Tökölyi, J., Schmidt, J. & Barta, Z. (2014) Climate and mammalian life histories. *Biological Journal of the Linnean Society*, **111**, 719-736.
- Vasseur, D.A., DeLong, J.P., Gilbertet, B. *et al.* (2014) Increased temperature variation poses a greater risk to species than climate warming. *Proceedings of the Royal Society of London Series B - Biological Sciences*, **281**, 20132612.

1  
2 630 Végvári, Z., Bókony, V., Barta, Z. & Kovács, G. (2010) Life history predicts advancement of avian  
3  
4 631 spring migration in response to climate change. *Global Change Biology*, **16**, 1-11.  
5  
6 632 Vincze, O., Székely, T., Küpper, C. *et al.* (2013) Local environment but not genetic differentiation  
7  
8 633 influences biparental care in ten plover populations. *PLoS ONE*, **8**, e60998.  
9  
10  
11 634 Walther, G.R., Post, E., Convey, P. *et al.* (2002) Ecological responses to recent climate change. *Nature*,  
12  
13 **416**, 389-395.  
14  
15  
16 636 Webb, D.R. (1987) Thermal tolerance of avian embryos: a review. *Condor*, **89**, 874–898.  
17  
18 637 Webb, J.N., Székely, T., Houston, A.I. & McNamara, J.M. (2002) A theoretical analysis of the energetic  
19  
20 costs and consequences of parental care decisions. *Philosophical Transactions of the Royal*  
21 638  
22 *Society B: Biological Sciences*, **357**, 331-340.  
23 639  
24  
25 640 Wilson, E. (1975) *Sociobiology: The New Synthesis*. Harvard University Press, Cambridge.  
26  
27  
28 641 Winkler, D.W., Dunn, P.O. & McCulloch, C.E. (2002) Predicting the effects of climate change on avian  
29  
30 642 life-history traits. *Proceedings of the National Academy of Sciences of the United States of*  
31  
32 *America*, **99**, 13595-13599.  
33 643  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1** Male incubation (binary response variable) in different plover species and populations (n = 5591 individuals). Mixed effects models.  $\chi^2$  values, degrees of freedom (df) and probability (p) of likelihood ratio tests are given.

		$\chi^2$ (df)	P
Model 1	<b>Fixed term</b>		
	Time period	1017.95 (9)	< 0.0001
	<b>Random terms</b>		
	Species	9.65 (1)	0.0019
	Population	44.91 (1)	< 0.0001
	Nest ID	0.00 (1)	1.000
Model 2	<b>Fixed term</b>		
	Time period	64.58 (9)	< 0.0001
	<b>Random terms</b>		
	Population	38.26 (1)	< 0.0001
	Species x time period	36.87 (1)	< 0.0001
Model 3	<b>Fixed term</b>		
	Time period	176.43 (9)	< 0.0001
	<b>Random terms</b>		
	Species	11.37 (1)	0.0007
	Population x time period	85.05 (1)	< 0.0001
Model 4	<b>Fixed terms</b>		
	Time period	1216.20 (63)	< 0.0001
	Mean temperature (°C)		
	Interaction with time period	84.42 (18)	< 0.0001
	Quadratic effect	32.03 (10)	0.0004
	Between-year temperature variation (°C)		
	Interaction with time period	15.23 (18)	0.6462
	Quadratic effect	2.82 (1)	0.0929
	Linear effect	7.34 (1)	0.0067
	Within-season temperature variation (°C)		
	Interaction with time period	70.81 (18)	< 0.0001
	Quadratic effect	33.68 (10)	0.0002
	<b>Random terms</b>		
	Species	14.07 (1)	0.0002
	Population	0.05 (1)	0.8298
	Nest identity	0.00 (1)	1.0000

**Footnote:** Main effects were tested by removing the main term and all its interactions with other variables. Interaction terms were tested by removing the interaction from full model and comparing the resulting model to the original. Quadratic terms were tested



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

650 by replacing polynomial with linear terms, and comparing the resulting model to the  
651 original.

For Peer Review

Figure legends


**Figure 1.** Male share of nest attendance estimated from capture data (% capture) in relation to male share of nest attendance as obtained from behavioural observations (% behaviour). Each point represents a 2-hour time period. Dashed lines represent equal estimates by the two methods. Statistics on each panel show the results of a least-squares regression weighted by the number of captures in each time period.  $R^2$  represents the coefficient of determination.

**Figure 2.** Male share of nest attendance (%) calculated from capture data in 36 populations. Each species is plotted on different panel, except Kentish plover and snowy plover which are shown on 3 and 2 panels, respectively. Each line represents a population. Legends refer to location numbers on the map (see Table S1 for population names and exact coordinates).

**Figure 3** Daily changes in predicted probability of male care (i.e. capture) in relation to (a) mean temperature, (b) between-year variation and (c) within-season variation. Each panel shows a different time period (see panel title for time period). Dashed lines represent 95% confidence intervals. Predictions are based on minimal model 4 from which the non-significant interaction and quadratic terms for between-year variation were removed (Table 1).

**Figure 4.** Predicted probability of male care (i.e. male capture) throughout the day under different climate scenarios. Each panel shows a climate scenario where the candidate temperature variable (i.e. shown by the main title of each sub-graph) takes three values (i.e., 2.5% quantile, median, 97.5% quantile), while the other two temperature variables are set to their median. Predictions are based on minimal model 4 from which the non-significant interaction and quadratic terms for between-year variation were removed (Table 1).

674 **Figure 1.**





675 **Figure 2.**

Figure 3 (a)



677 Figure 3 (b)




Figure 3 (c)






Figure 4.





SUPPORTING INFORMATION

**Figure S1.** Schematic presentation of how the effect of each variable was tested. All derived models (i.e. shown by arrowheads) were compared to the model from which they originate (i.e. shown by



the base of the arrows). Models were compared using likelihood ratio statistics.

**Table S1** Summary of parental care data from different populations of plovers *Charadrius spp.* Sexing method refers to molecular sexing (M), plumage and/or other morphometric measurements or behaviour based (P). Numbers in square brackets in the Population column refer to the localities on the map in Figure 2. See Appendix S1 for references cited here.

Species	Population	Coordinates	Years of data collection	Number of captures	% male captures	Sexing method	References
<i>C. alexandrinus</i>	Maio Island (Cape Verde) [15]	15°09'N, 23°13'W	2007-2010	244	40.57	M, P	Székely T., A.A. Tico & A. Kosztolányi unpubl data
<i>C. alexandrinus</i>	Farasan Islands (Saudi Arabia) [26]	16°48'N, 41°53'E	2008-2009, 2011	45	35.56	P	AlRashidi <i>et al.</i> , 2011
<i>C. alexandrinus</i>	Al Wathba Wetland (United Arab Emirates) [27]	24°16'N, 54°36'E	2005-2006	175	48.00	P	Kosztolányi <i>et al.</i> , 2009, AlRashidi <i>et al.</i> , 2010
<i>C. alexandrinus</i>	Tuzla Lake (Turkey) [25]	36°42'N, 35°03'E	1996-2000, 2004	604	46.19	P	Kosztolányi & Székely 2002
<i>C. alexandrinus</i>	Fuente de Piedra Lake (Spain) [16]	37°06'N, 04°45'W	1991-1996	174	49.43	P	Fraga & Amat (1996), Amat & Masero 2004
<i>C. alexandrinus</i>	Bohai Bay (China) [28]	39°05'N, 118°12'E	2012	38	31.58	M, P	Que, P. & Y. Liu unpubl. data
<i>C. alexandrinus</i>	Llobregat Delta (Spain) [17]	41°18'N, 02°08'E	1994-1995, 1998, 2000-2008	173	41.71	P	Figuerola & Cerdà 1998
<i>C. alexandrinus</i>	Lagoon of Venice and Po Delta (Italy) [23]	45°10'N, 12°24'E	1993-1995	157	45.86	P	Serra, L. unpubl. data
<i>C. alexandrinus</i>	Great Hungarian Plain (Hungary) [24]	46°40'N, 19°10'E	1988-1994	186	39.25	P	Székely & Lessells 1993, Székely <i>et al.</i> , 1994
<i>C. alexandrinus</i>	Schleswig-Holstein (Germany) [21]	54°45'N, 08°01'E	1989-1998, 2001-2002, 2004-2005	530	44.34	P	Schulz, R. unpubl. data
<i>C. alexandrinus</i>	Falsterbo Peninsula (Sweden) [22]	55°15'N, 12°34'E	1981-1988, 1990-1991, 1993-1994, 1996	44	47.73	P	Jönsson, P. unpubl. data
<i>C. falklandicus</i>	Sea Lion Island (Falklands) [13]	51°41'S, 59°10'W	2005-2008	63	42.86	M, P	St Clair <i>et al.</i> , 2010a
<i>C. falklandicus</i>	Peninsula Valdés (Argentina) [12]	42°30'S, 63°56'W	2006-2007	62	36.51	M, P	García-Peña 2009
<i>C. marginatus</i>	Cape Peninsula (South Africa) [18]	34°08'S, 18°20'E	1999-2003	162	32.72	P	Lloyd, P. unpubl. data
<i>C. marginatus</i>	Lake Tsimanampetsotsa (Madagascar) [20]	24°48'S, 43°49'E	2005-2006, 2011-2012	41	43.90	M	Zefania, S, J. Parra & T. Székely unpubl. data

<i>C. marginatus</i>	Andavadoaka saltmarsh (Madagascar) [19]	22°04'S, 43°14'E	2010-2012	48	43.75	M	Zefania, S, J. Parra & T. Székely unpubl. data
<i>C. melodus</i>	Saskatchewan Coteau (Canada) [1]	49°44'N, 105°23'W	2002-2006	139	49.64	P	Cohen & Gratto-Trevor 2011; Gratto-Trevor 2011
<i>C. melodus</i>	Saskatchewan Diefenbaker (Canada) [2]	50°43'N, 107°30'W	2002-2007	268	49.44	P	Cohen & Gratto-Trevor 2011; Gratto-Trevor 2011
<i>C. melodus</i>	Saskatchewan Quill (Canada) [3]	51°55'N, 104°22'W	2002-2006	176	49.15	P	Cohen & Gratto-Trevor 2011; Gratto-Trevor 2011
<i>C. modestus</i>	Sea Lion Island (Falklands) [13]	51°41'S, 59°10'W	2005-2008	99	55.56	M, P	St Clair <i>et al.</i> , 2010a, St Clair <i>et al.</i> , 2010b
<i>C. nivosus</i>	Texcoco (Mexico) [11]	19°30'N, 98°29'W	2009-2012	57	21.05	P	DeSucre-Medrano, A. E. & S. Gomez del Angel unpubl. data
<i>C. nivosus</i>	Nayarit (Mexico) [10]	22°16'N, 105°12'W	2010-2012	44	40.91	P	Villar, C. & J. Cavitt unpubl. data
<i>C. nivosus</i>	Ceuta Bay (Mexico) [9]	23°54'N, 106°57'W	2006-2012	451	48.12	P	Küpper, C. & M. Cruz-López unpubl. data
<i>C. nivosus</i>	Florida (USA) [4]	29°44'N, 85°06'W	2008-2010	300	10.33		Pruner, R. unpubl. data
<i>C. nivosus</i>	San Quintin Bay (Mexico) [7]	30°40'N, 116°0'W	2012	45	19.57	P	Galindo-Espinosa, D. unpubl. data
<i>C. nivosus</i>	Texas (USA) [8]	33°12'N, 102°30'W	1999-2000, 2008-2009	127	33.86	P, M	Saalfeld <i>et al.</i> , 2011
<i>C. nivosus</i>	Monterey Bay (USA) [6]	36°45'N, 121°25'W	1984-1999	581	18.93	P	Warriner <i>et al.</i> , 1986, Stenzel <i>et al.</i> , 2011
<i>C. nivosus</i>	Great Salt Lake (USA) [5]	41°41'N, 112°55'W	2007-2010	80	22.50	P	Cavitt <i>et al.</i> , 2008, Hall & Cavitt 2012
<i>C. pecuarius</i>	Lake Tsimanampetsotsa (Madagascar) [20]	24°48'S, 43°49'E	2005, 2007, 2012	37	43.24	M	Zefania, S., J. Parra & T. Székely unpubl. data
<i>C. pecuarius</i>	Andavadoaka saltmarsh (Madagascar) [19]	22°04'S, 43°14'E	2010, 2012	118	49.15	M	Zefania, S., J. Parra & T. Székely unpubl. data
<i>C. peronii</i>	Prachuap Khiri Khan (Thailand) [29]	12°00'N, 99°53'E	2004-2005	65	46.97	P	Yasué & Dearden 2006a,b, 2007a,b
<i>C. ruficapillus</i>	Altona Saltworks (Australia) [30]	37°53'S, 144°47'E	2008-2012	71	36.62	P	Lomas <i>et al.</i> , 2014, Weston, M.A. unpubl. data
<i>C. sanctaehelenae</i>	St. Helena Island (St. Helena) [14]	15°58'S, 05°43'W	2004, 2007-2009	48	41.67	P, M	Burns <i>et al.</i> , 2013
<i>C. thoracicus</i>	Lake Tsimanampetsotsa (Madagascar) [20]	24°48'S, 43°49'E	2004-2009, 2011-2012	93	31.18	M	Zefania, S, J. Parra & T. Székely unpubl. data
<i>C. thoracicus</i>	Andavadoaka saltmarsh (Madagascar) [19]	22°04'S, 43°14'E	2010	19	31.58	M	Zefania, S, J. Parra & T. Székely unpubl. data
<i>C. wilsonia</i>	Ceuta Bay (Mexico) [9]	23°54'N, 106°57'W	2009, 2012	27	37.04	P, M	Küpper, C. & M. Cruz-López unpubl. data

**Footnote:** Molecular sexing markers: P2P8, Z-002B and Calnex-31 (Griffiths *et al.*, 1998, Dawson 2007, Küpper *et al.*, 2007)

**Table S2.** Sensitivity analyses for the length of the time period on which the calculation of the three climate variables was based on.

Model 4	Fixed terms	5 years		10 years		15 years	
	Time period	1214.40 (63)	< 0.0001	1219.2 (63)	< 0.0001	1217.70 (63)	< 0.0001
	<u>Mean temperature (°C)</u>						
	Interaction with time period	79.96 (18)	< 0.0001	77.28 (18)	< 0.0001	85.49 (18)	< 0.0001
	Quadratic effect	32.93 (10)	0.0003	32.76 (10)	0.0003	33.26 (10)	0.0002
	<u>Between-season temperature variation (°C)</u>						
	Interaction with time period	37.86 (18)	0.0040	23.35 (18)	0.1777	15.76 (18)	0.6091
	Quadratic effect	16.32 (10) *	0.0907	1.52 (1)	0.2173	1.22 (1)	0.2690
	Linear effect	6.19 (2) *	0.0452	4.67 (1)	0.0307	6.03 (1)	0.0140
	<u>Within-season temperature variation (°C)</u>						
	Interaction with time period	40.71 (18)	0.0017	32.57 (18)	0.0188	70.65 (18)	< 0.0001
	Quadratic effect	23.97 (10)	0.0077	32.76 (10)	0.0003	29.36 (10)	0.0011
	<b>Random terms</b>						
	Species	16.40 (1)	< 0.0001	14.49 (1)	0.0001	14.46 (1)	0.0001
	Population	0.22 (1)	0.6367	0.60 (1)	0.4400	0.07 (1)	0.7919
	Nest identity	0.00 (1)	0.9984	0.00 (1)	1.0000	0.00 (1)	1.0000

\*Interaction with time period retained in model (similarly to the other two temperature variables) due to its significant effect.

**Appendix S1.** Supplementary references for Table S1

AlRashidi M, Kosztolányi A, Küpper C, Cuthill IC, Javed S, Székely T (2010) The influence of a hot environment on parental cooperation of a ground-nesting shorebird, the Kentish plover *Charadrius alexandrinus*. *Frontiers in Zoology*, **7**, 1.

AlRashidi M, Kosztolányi A, Shobrak M, Küpper C, Székely T (2011) Parental cooperation in an extreme hot environment: natural behaviour and experimental evidence. *Animal Behaviour*, **82**, 235–243.

Amat JA, Masero JA (2004) How Kentish plovers, *Charadrius alexandrinus*, cope with heat stress during incubation. *Behavioral Ecology and Sociobiology*, **56**, 26-33.

Burns F, McCulloch N, dos Remedios N, Bolton M, Székely T (2013) Sex differences in incubation behaviour but not mortality risk in a threatened shorebird. *Ibis*, **155**, 877-880.

Cavitt JF, Cole L, Stone K (2008) Breeding Productivity of Shorebirds and Colonial Waterbirds at Bear River Migratory Bird Refuge, Utah. Technical Report to US Fish and Wildlife Service, Bear River Migratory Bird Refuge, Brigham City, UT.

Cohen JB, Gratto-Trevor CL (2011) Survival, site fidelity, and population dynamics of Piping Plovers in Saskatchewan, *Journal of Field Ornithology*, **82**, 372-394.

Dawson DA (2007) Genomic analysis of passerine birds using conserved microsatellite loci. PhD Thesis, University of Sheffield, UK.

Fraga RM, Amat JA (1996) Biología de la reproducción de una población de Chorlitejos Patinegros (*Charadrius alexandrinus*) en una laguna salada interior. *Ardeola*, **43**, 69-85.

García-Peña GE (2009) Phylogenetic comparative analyses of breeding systems and life-history strategies in shorebirds. Publisher: University of Bath.

Gratto-Trevor CL (2011) Ageing and sexing the Piping Plover *Charadrius melodus*. *Wader Study Group Bulletin*, **118**, 118–122.

- Griffiths R, MC Double, K Orr, Dawson RJG (1998) A DNA test to sex most birds. *Mol. Ecol.* **7**: 1071–1075. Figuerola J, Cerdà F (1998) Evolució i conservació de la població de corriol camanegre (*Charadrius alexandrinus*) del delta del Llobregat. *Spartina*, **3**, 161-169.
- Hall L, Cavitt JF (2012) Comparative study of trapping methods for ground-nesting shorebirds. *Waterbirds* **35**, 342-346.
- Kosztolányi A, Székely T (2002) Using a transponder system to monitor incubation routines of snowy plovers. *Journal of Field Ornithology*, **73**, 199-205.
- Kosztolányi A, Javed S, Küpper C, Cuthill IC, Al Shamsi A, Székely T (2009) Breeding ecology of Kentish Plover *Charadrius alexandrinus* in an extremely hot environment. *Bird Study*, **56**, 244-252.
- Küpper C, Horsburgh GJ, Dawson DA, Ffrench-Constant R, Székely T, Burke T (2007) Characterisation of 36 polymorphic microsatellite loci in the Kentish Plover (*Charadrius alexandrinus*) including two sex-linked loci and their amplification in four other *Charadrius* species. *Molecular Ecology Notes*, **7**, 35-39.
- Lomas SC, Whisson DA, Maguire GS, Tan LX, Guay P-J, Weston MA (2014) The influence of cover on nesting Red-capped Plovers: a trade-off between thermoregulation and predation risk? *Victorian Naturalist*, **131**, 115-127.
- Saalfeld ST, Conway WC, Haukos DA, Johnson WP (2011) Nest success of snowy plovers (*Charadrius nivosus*) in the Southern High Plains of Texas. *Waterbirds*, **34**, 389-399.
- Székely T, Lessells CM (1993) Mate change by Kentish plovers *Charadrius alexandrinus*. *Ornis Scandinavica*, **24**, 317-322.
- Székely T, Karsai I, Williams TD (1994) Determination of clutch size in the Kentish Plover *Charadrius alexandrinus*. *Ibis*, **136**, 341-348.
- St Clair JJH, Küpper C, Herrmann P, Woods RW, Székely T (2010a) Unusual incubation sex-roles in the Rufous-chested Dotterel *Charadrius modestus*. *Ibis*, **152**, 402



1  
2  
3 St Clair JJH, Herrmann P, Woods RW, Székely T (2010b) Female-biased incubation and strong diel sex-  
4  
5 roles in the Two-banded Plover *Charadrius falklandicus*. *Journal of Ornithology*, **151**, 811-816  
6  
7 Stenzel LE, et al. (2011) Male-skewed adult sex ratio, survival, mating opportunity and annual  
8  
9 productivity in the Snowy Plover *Charadrius alexandrinus*. *Ibis*, **153**, 312-322.  
10  
11 Warriner JS, Warriner JC, Page GW, Stenzel LE (1986) Mating system and reproductive success of a  
12  
13 small population of polygamous Snowy Plovers. *Wilson Bulletin*, **98**, 15-37.  
14  
15  
16 Yasué, M. & Dearden, P. (2006a). The effects of heat stress, predation risk and parental investment on  
17  
18 Malaysian plover nest return times after a human disturbance. *Biological Conservation*, **132**, 472-  
19  
20 480.  
21  
22  
23 Yasué M, Dearden P (2006b) Simultaneous biparental incubation of two nests by a pair of Malaysian  
24  
25 plovers. *Wader Study Group Bulletin*, **109**, 121 – 122.  
26  
27  
28 Yasué M, Dearden P (2007a) Constraints on successive clutching behaviour of sedentary Malaysian  
29  
30 plovers breeding in a tropical environment. *Ardea*, **96**, 59-72.  
31  
32 Yasué M, Dearden P (2007b) Sex-roles of Malaysian plovers *Charadrius peronii* during territory  
33  
34 acquisition, incubation and brood care. *Journal of Ethology*, **26**, 99-112.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60